

GENERALIZATIONS OF THE ROWLAND THEOREM

VLADIMIR SHEVELEV

ABSTRACT. We prove the theorems which are equivalent to the Rowland's results such that a new form of them allows to consider some generalizations. In particular, we give generators of primes more than a fixed prime.

1. INTRODUCTION

In 2008, Rowland [1], using very elementary tools, discovered a very interesting fact.

Theorem 1. *Let $a(1) = 7$ and, for $n \geq 2$,*

$$a(n) = a(n-1) + \gcd(n, a(n-1)).$$

Then, for every $n \geq 2$, the difference $a(n) - a(n-1)$ is 1 or prime.

He also mentioned that, in particular, the following similar theorem is true (it also follows from his proof).

Theorem 2. *Let $b(3) = 6$ and, for $n \geq 4$,*

$$b(n) = b(n-1) + \gcd(n, b(n-1)).$$

Then, for every $n \geq 2$, the difference $b(n) - b(n-1)$ is 1 or prime.

In spite of the nearness of Theorem 1 and 2, there is an essential distinction: $\limsup a(n)/n = 3$, while $\limsup b(n)/n = 2$. An infinite sets of the initial conditions for which one of Theorems 1,2 is true we considered in [2].

A generalization of Theorems 1,2 could appear if to become free from the hard dependence on the value of $\gcd$.

It is important that the Rowland's proof allows to write his results in such a form. The following theorems are equivalent to Theorems 1,2.

Theorem 1a. *Let $a(1) = 7$ and, for $n \geq 2$,*

$$a(n) = \begin{cases} a(n-1) + 1, & \text{if } \gcd(n, a(n-1)) = 1 \\ 3n, & \text{otherwise} \end{cases}.$$

Then, for every $n \geq 2$, the difference $a(n) - a(n-1)$ is 1 or prime.

Theorem 2a. *Let $b(3) = 6$ and, for $n \geq 4$,*

$$b(n) = \begin{cases} b(n-1) + 1, & \text{if } \gcd(n, b(n-1)) = 1 \\ 2n, & \text{otherwise} \end{cases}.$$

Then, for every $n \geq 4$, the difference $b(n) - b(n-1)$ is 1 or prime.

2. DIRECT PROOF OF THEOREMS 1a, 2a WITH GENERAL INITIALS

Excepting a trivial case when $a(n) = 1$ identically, the sequences which are defined by Theorems 1a and 2a contain the values in which $a(n) = 3n$ or $a(n) = 2n$ correspondingly. Therefore the general initial conditions for them have the form $a(n_1) = 3n_1$ or $a(n_1) = 2n_1$ correspondingly. Below we give a direct and simple proofs of Theorems 1a, 2a with general initial conditions. Our proof is again based on Rowland idea, but in some another form.

Theorem 1b. Let, for $n_1 \geq 2$, $a(n_1) = 3n_1$ and, for $n \geq n_1 + 1$,

$$(2.1) \quad a(n) = \begin{cases} a(n-1) + 1, & \text{if } \gcd(n, a(n-1)) = 1, \\ 3n, & \text{otherwise} \end{cases}.$$

Then, for every $n \geq n_1 + 1$, the difference $a(n) - a(n-1)$ is 1 or prime.

Theorem 2b. Let, for $n_1 \geq 3$, $b(n_1) = 2n_1$ and, for $n \geq n_1 + 1$,

$$(2.2) \quad b(n) = \begin{cases} b(n-1) + 1, & \text{if } \gcd(n, b(n-1)) = 1, \\ 2n, & \text{otherwise} \end{cases}.$$

Then, for every $n \geq n_1 + 1$, the difference $b(n) - b(n-1)$ is 1 or prime.

Proof of Theorem 2b. Let k be the smallest positive integer such that

$$(2.3) \quad \gcd(n_1 + k, 2n_1 + k - 1) = d > 1.$$

Then from (2.2) and 2.3) we have

$$(2.4) \quad \begin{aligned} b(n_1) &= 2n_1, \\ b(n_1 + 1) &= 2n_1 + 1, \\ &\dots \\ b(n_1 + k - 1) &= 2n_1 + k - 1, \\ b(n_1 + k) &= 2(n_1 + k). \end{aligned}$$

Besides, by (2.3), $\gcd(n_1 - 1, n_1 + k) = d$ and, therefore,

$$(2.5) \quad \gcd(n_1 - 1, k + 1) = d.$$

Thus some prime divisor P of $n_1 - 1$ divides $k + 1$, i.e. $k + 1 \geq P$. All the more,

$$(2.6) \quad k + 1 \geq p,$$

where p is the smallest prime divisor of $n_1 - 1$ (by the condition, $n_1 - 1 \geq 2$).

On the other hand, inside (2.4) there is a row

$$(2.7) \quad b(n_1 + p - 2) = 2n_1 + p - 2.$$

Since $\gcd(2n_1 + p - 2, n_1 + p - 1) \geq p$, then, by the definition of k , we have $k \leq p - 1$ and, in view of (2.6),

$$(2.8) \quad k + 1 = p.$$

Now from (2.4) we find

$$\begin{aligned} b(n_1 + k) - b(n_1 + k - 1) &= k + 1 = p, \\ b(n_1 + k) &= 2(n_1 + k) \end{aligned}$$

and, by the evident induction, we are done. ■

Proof of Theorem 1b is the same if to replace (2.4) by

$$a(n_1 + k - 1) = 3n_1 + k - 1,$$

$$a(n_1 + k) = 3(n_1 + k),$$

(2.5) by

$$\gcd(2n_1 - 1, k + 1) = d$$

and (2.7) by

$$a(n_1 + \frac{p-1}{2} - 1) = 3n_1 + \frac{p-1}{2} - 1,$$

where now p is the smallest prime divisor of $2n_1 - 1 \geq 3$, such that $2k + 1 = p$. ■

3. ALGORITHMS OF FAST CALCULATION OF NONTRIVIAL INCREMENTS IN SEQUENCES OF THEOREMS 1b, 2b

Sequences of the considered type contain too many points of trivial 1-increments. Nevertheless, it is interesting to see quicker what primes these sequences contain. Therefore, the following problem is actual from the computation point of view : to accelerate this algorithm for receiving of primes by the omitting of the trivial increments. Using induction, from the proofs of theorems 1b, 2b we discover a sense of these primes by the following algorithms.

For integer $n \geq 2$, denote n^* the least prime divisor of n . We start with an algorithm of fast calculation of the Rowland's primes which are obtained by Theorem 1.

Algorithm 1. *Put*

$$N_1 = 4, \quad p_1 = 5$$

and, for $i \geq 1$, put

$$N_{i+1} = N_i + p_i - 1, \quad p_{i+1} = (N_{i+1} + 1)^*.$$

Then sequence $\{p_i\}_{i \geq 1}$ is the sequence of nontrivial increments of the Rowland sequence, which is defined by Theorem 1.

By this algorithm, we consecutively find

$$p_1 = 5, \quad N_2 = 8, \quad p_2 = 9^* = 3, \quad N_3 = 10, \quad p_3 = 11, \quad N_4 = 20, \quad p_4 = 3, \dots$$

Using formulas of Algorithm 1, it is easy to obtain the following recursion for p_n with the automatically produced initial term:

$$(3.1) \quad p_n = (6 - n + \sum_{i=1}^{n-1} p_i)^*.$$

Example 1. We have consecutively (cf. sequence A137613 in [3]):

$$\begin{aligned} p_1 &= (6 - 1)^* = 5, \\ p_2 &= (6 - 2 + 5)^* = 3, \\ p_3 &= (6 - 3 + 5 + 3)^* = 11, \\ p_4 &= (6 - 4 + 5 + 3 + 11)^* = 3, \\ p_5 &= (6 - 5 + 5 + 3 + 11 + 3)^* = 23, \end{aligned}$$

etc.

Note that, beginning with $n \geq 3$, the Rowland's sequence could be defined by the initial condition $a(3) = 9$. Therefore, the following algorithm is a generalization of Algorithm 1.

Algorithm 2. Consider sequence of Theorem 1b with initial condition $a(n_1) = 3n_1$. Put

$$N_1 = 2(n_1 - 1), \quad p_1 = (N_1 + 1)^*$$

and, for $i \geq 1$, put

$$N_{i+1} = N_i + p_i - 1, \quad p_{i+1} = (N_{i+1} + 1)^*.$$

Then sequence $\{p_i\}_{i \geq 1}$ is the sequence of nontrivial increments of the sequence, which is defined by Theorem 1b.

E.g., for $n_1 = 4$, by this algorithm, we consecutively find

$$N_1 = 6, \quad p_1 = 7, \quad N_2 = 12, \quad p_2 = 13, \quad N_3 = 24, \quad p_3 = 5, \quad N_4 = 28, \quad p_4 = 29, \dots$$

Using formulas of Algorithm 2, we obtain the following recursion for p_n with the automatically produced initial term:

$$(3.2) \quad p_n = (2n_1 - n + \sum_{i=1}^{n-1} p_i)^*,$$

such that, for $n_1 = 3$, $n \geq 3$, we have (3.1).

Example 2. *In case of $n_1 = 4$, we have consecutively:*

$$\begin{aligned} p_1 &= (8 - 1)^* = 7, \\ p_2 &= (8 - 2 + 7)^* = 13, \\ p_3 &= (8 - 3 + 7 + 13)^* = 5, \\ p_4 &= (8 - 4 + 7 + 13 + 5)^* = 29, \end{aligned}$$

etc.

Finally, we give the corresponding algorithm for sequences which are defined by Theorem 2b.

Algorithm 3. *Consider sequence of Theorem 2b with initial condition $a(n_1) = 2n_1$. Put*

$$N_1 = n_1 - 2, \quad p_1 = (N_1 + 1)^*$$

and, for $i \geq 1$, put

$$N_{i+1} = N_i + p_i - 1, \quad p_{i+1} = (N_{i+1} + 1)^*.$$

Then sequence $\{p_i\}_{i \geq 1}$ is the sequence of nontrivial increments of the sequence, which is defined by Theorem 2b.

E.g., for $n_1 = 5$, by this algorithm, we consecutively find

$$N_1 = 3, \quad p_1 = 2, \quad N_2 = 4, \quad p_2 = 5, \quad N_3 = 8, \quad p_3 = 3, \quad N_4 = 10, \quad p_4 = 11, \dots$$

Here we obtain the following recursion for p_n :

$$(3.3) \quad p_n = (n_1 - n + \sum_{i=1}^{n-1} p_i)^*.$$

Example 3. *In case of $n_1 = 5$, we have consecutively:*

$$\begin{aligned} p_1 &= (5 - 1)^* = 2, \\ p_2 &= (5 - 2 + 2)^* = 5, \\ p_3 &= (5 - 3 + 2 + 5)^* = 3, \\ p_4 &= (5 - 4 + 2 + 5 + 3)^* = 11, \end{aligned}$$

etc.

4. ν - GENERALIZATIONS: OTHER GENERATORS OF PRIMES

Theorem 3. *Let ν be a positive even integer. Let, for $n_1 \geq 2$, $c(n_1) = 3n_1 + \nu$ and, for $n \geq n_1 + 1$,*

$$(4.1) \quad c(n) = \begin{cases} 3n + \nu, & \text{if } \gcd(n, c(n-1)) > 1 \text{ and } \rho \geq \frac{\nu-2}{2}, \\ c(n-1) + 1, & \text{otherwise} \end{cases}$$

where, for $n > n_1$,

$$(4.2) \quad \rho = \rho(n) := n - \max\{l < n : c(l) = 3l + \nu\}.$$

Then, for every $n \geq n_1 + 1$, the difference $c(n) - c(n-1) - \nu$ is 1 or ν or prime.

Note that, condition (4.2) means that in this sequence the distance between the consecutive nontrivial increments (i.e. the increments which are different from 1) is not less than $\frac{\nu-2}{2}$.

Proof. Let k be the smallest positive integer not less than $\frac{\nu-2}{2}$ such that

$$(4.3) \quad \gcd(n_1 + k, 3n_1 + k - 1 + \nu) = d > 1.$$

Then from (4.1) and 4.3) we have

$$(4.4) \quad \begin{aligned} c(n_1) &= 3n_1 + \nu, \\ c(n_1 + 1) &= 3n_1 + 1 + \nu, \\ &\dots \\ c(n_1 + k - 1) &= 3n_1 + k - 1 + \nu, \\ c(n_1 + k) &= 3(n_1 + k) + \nu. \end{aligned}$$

Besides, by (4.3), $\gcd(2n_1 + \nu - 1, n_1 + k) = d > 1$ and, therefore,

$$(4.5) \quad \gcd(2n_1 + \nu - 1, 2k - \nu + 1) = d > 1.$$

Thus some prime divisor P of $2n_1 + \nu - 1$ divides $2k - \nu + 1$, i.e. $2k - \nu + 1 \geq P$. All the more,

$$(4.6) \quad 2k - \nu + 1 \geq p,$$

where p is the smallest prime divisor of $2n_1 + \nu - 1$ (by the condition, ν is even, therefore p is odd). Note that (4.6) shows that the condition $k \geq \frac{\nu-2}{2}$ is necessary.

On the other hand, inside (2.4) there is a row

$$(4.7) \quad c(n_1 + \frac{p + \nu - 3}{2}) = 3n_1 + \frac{p + \nu - 3}{2} + \nu = \frac{1}{2}(6n_1 + 3\nu - 3 + p).$$

Since $n_1 + \frac{p + \nu - 1}{2} = \frac{1}{2}(2n_1 + \nu - 1 + p)$, then we see that

$$\gcd(c(n_1 + \frac{p + \nu - 3}{2}), n_1 + \frac{p + \nu - 1}{2}) \geq p,$$

and, by the definition of k , we have $k \leq \frac{p+\nu-1}{2}$, and, in view of (2.6), we conclude that

$$(4.8) \quad 2k - \nu + 1 = p.$$

Now from (2.4) we find

$$c(n_1 + k) - c(n_1 + k - 1) - \nu = 2k + 1 - \nu = p,$$

and, for $n_2 := n_1 + k$,

$$c(n_2) = 3(n_2) + \nu$$

and, by the evident induction, we are done. ■

The following theorem is proved quite analogously.

Theorem 4. *Let ν be a positive integer. Let, for $n_1 \geq 2$, $c(n_1) = 2n_1 + \nu$ and, for $n \geq n_1 + 1$,*

$$(4.9) \quad c(n) = \begin{cases} 2n + \nu, & \text{if } \gcd(n, c(n-1)) > 1 \text{ and } \rho \geq \nu + 1, \\ c(n-1) + 1, & \text{otherwise} \end{cases}$$

where, for $n > n_1$,

$$(4.10) \quad \rho = \rho(n) := n - \max\{l < n : c(l) = 2l + \nu\}.$$

Then, for every $n \geq n_1 + 1$, the difference $c(n) - c(n-1) - \nu$ is 1 - ν or prime.

5. ν - GENERALIZATIONS: GENERATORS OF PRIMES MORE THAN p_m

The following theorem gives generators of primes $p > p_m$, $m \geq 3$, where p_n is the n -th prime.

Theorem 5. *Let $m \geq 3$, $\nu \geq 0$ be even and not exceeding $p_m - 3$. Let, for $n_1 > \nu + 2$, $c(n_1) = 3n_1 - \nu$ and, for $n \geq n_1 + 1$,*

$$(5.1) \quad c(n) = \begin{cases} 3n - \nu, & \text{if } \gcd(n, c(n-1)) > 1 \text{ and } \gcd(n, \prod_{i=1}^m p_i) = 1, \\ c(n-1) + 1. & \text{otherwise} \end{cases}$$

Then, for every $n \geq n_1 + 1$, the enlarged on ν difference $c(n) - c(n-1)$ is $1 + \nu$ or prime more than p_m .

Proof is similar to proof of Theorem 3 with the replacing ν by $-\nu$. Let k be the smallest positive integer such that

$$\gcd(n_1 + k, \prod_{i=1}^m p_i) = 1, \text{ and}$$

$$(5.2) \quad \gcd(n_1 + k, 3n_1 + k - 1 + \nu) = d > 1.$$

Then, as in proof of Theorem 3, we conclude that

$$(5.3) \quad \gcd(2n_1 - \nu - 1, 2k + \nu + 1) = d > 1 \text{ and} \\ \gcd(2n_1 - \nu - 1, \prod_{i=1}^m p_i) = 1.$$

Thus if p is the smallest prime divisor of $2n_1 - \nu - 1$, then

$$(5.4) \quad p > p_m \geq \nu + 3,$$

and

$$(5.5) \quad 2k + \nu + 1 \geq p.$$

Note that the condition $n_1 > \nu + 2$ is necessary. Indeed, by (5.3), we have $2n_1 - \nu - 1 \geq p > p_m \geq \nu + 3$ and the inequality $n_1 > \nu + 2$ follows.

On the other hand, in view of (5.4) inside (2.4) with the replacing ν by $-\nu$ we find a row

$$(5.6) \quad c(n_1 + \frac{p - \nu - 3}{2}) = 3n_1 + \frac{p - \nu - 3}{2} - \nu = \frac{1}{2}(6n_1 - 3\nu - 3 + p).$$

Since $n_1 + \frac{p - \nu - 1}{2} = \frac{1}{2}(2n_1 - \nu - 1 + p)$, then we see that

$$\gcd(c(n_1 + \frac{p - \nu - 3}{2}), n_1 + \frac{p - \nu - 1}{2}) \geq p,$$

and, by the definition of k , we have $k \leq \frac{p - \nu - 1}{2}$, and, in view of (5.5), we conclude that

$$(5.7) \quad 2k + \nu + 1 = p.$$

Now we again obtain that

$$c(n_1 + k) - c(n_1 + k - 1) + \nu = 2k + 1 + \nu = p,$$

and, for $n_2 := n_1 + k$,

$$c(n_2) = 3(n_2) - \nu$$

and, by the evident induction, we are done. ■

The following theorem is proved quite analogously.

Theorem 6. *Let $m \geq 2$, ν be even not exceeding $p_m - 2$. Let, for $n_1 \geq 2\nu + 4$, $c(n_1) = 2n_1 - \nu$ and, for $n \geq n_1 + 1$,*

$$(5.8) \quad c(n) = \begin{cases} 2n - \nu, & \text{if } \gcd(n, c(n-1)) > 1 \text{ and } \gcd(n, \prod_{i=1}^m p_i) = 1, \\ c(n-1) + 1. & \text{otherwise} \end{cases}$$

Then, for every $n \geq n_1 + 1$, the enlarged on ν difference $c(n) - c(n - 1)$ is $1 + \nu$ or prime more than p_m .

Note that the conditions $\nu \leq p_m - 2$ and $n_1 \geq 2\nu + 4$ are necessary (cf. proof of Theorem 5).

6. EXAMPLES

1) An illustration to Theorem 3. $\nu = 4$, $n_1 = 6$. The first terms of the sequence are:

$$22, 23, 24, 31, 32, 33, 34, 35, 46, \dots$$

with the first nontrivial diminished on 4 increments $3, 7, 31, \dots$

2) An illustration to Theorem 4. $\nu = 5$, $n_1 = 15$. The first terms of the sequence are:

$$35, 36, 37, 38, 39, \dots$$

Note that $a(19) = 39$, since $19 - 15 < \nu + 1 = 6$.

The first nontrivial diminished on 5 increments are $19, 3, 7, 3, 67, \dots$

3) An illustration to Theorem 5 (cf. sequences A168143-A168144 in [3]). $p_7 = 17$, $\nu = 14$, $n_1 = 17$. The first terms of the sequence are:

$$37, 38, 43, 44, 45, 46, 55, 56, 57, 58, 73, \dots$$

with the first nontrivial enlarged on 14 increments $19, 23, 29, 43, 71, \dots$

Astonishingly that, probably, it is difficult to find a regularity in order to obtain a ν -generalization of algorithms of Section 3.

7. AN ADDITIONAL RESULT

An attempt to construct of similar sequences (in style of Theorems 3-6) with *increasing* nontrivial increments leads us to a surprising thing. More exactly, consider the following sequence. Let $a(3) = 6$ and, for $n \geq 4$,

$$(7.1) \quad c(n) = 2n, \text{ if } \gcd(n, c(n-1)) > 1$$

and for $m < n$,

$$\gcd(m, c(m-1)) < \gcd(n, c(n-1)),$$

and, otherwise,

$$c(n) = c(n-1) + 1.$$

Theorem 7. *The nontrivial increments of $\{c(n)\}$ form the sequence*

$$\{2^n + 1\}_{n \geq 1}.$$

Firstly, we prove a lemma.

Lemma 1. *For $j = 1, 2, \dots, n-1$, we have*

$$\gcd(n+j, 2n+j-1) \leq n/2.$$

Proof. Indeed, evidently

$$\gcd(n+j, 2n+j-1) = \gcd(n+j, n-1) = \gcd(j+1, n-1),$$

and the lemma follows. ■

Proof of Theorem 7. We use induction with the base $c(3) = 6$, $c(4) = 8$. We see that to obtain $c(4) = 8$ from $c(3) = 6$ we add 1 to the value of argument 3 and 0 to the term 6 of the sequence; after that we obtain $c(4) = c(3) + \gcd(3+1, c(3)+0) = 6 + \gcd(4, 6) = 8$. Suppose that in case of $c(n) = 2n$ we add $n-2$ to the value of argument n and $n-3$ to the term $c(n)$. Then we have

$$c(2n-2) = 3n-3 + \gcd(2n-2, 3n-3) = 4n-4.$$

By the same way, putting $2n-2 = m$, we have

$$c(2m-2) = 3m-3 + \gcd(2m-2, 3m-3) = 4m-4.$$

Using Lemma 1, we see that

$$\gcd(m+j, 2m+j-1) < m/2 = n-1,$$

i.e. the value of gcd of the precede step. Now, by the induction, we conclude that the numbers $2, 2 \cdot 2 - 1 = 3, 2 \cdot 3 - 1 = 5, \dots, 2^k + 1, \dots$ are the smallest values of $\gcd(n, c(n-1))$, to which do not precede any non-smaller values of gcd. ■

Close results we could prove, putting, e.g., $c(7) = 12$ (here we obtain numbers 4, 7, 13, 25, ...) or $c(5) = 12$ with the replacing in (6.1) $c(n) = 2n$ by $c(n) = 3n$ (here we obtain numbers 6, 11, 21, 41, ...) etc.

REFERENCES

- [1] . E. S. Rowland, *A natural prime-generating recurrence* J.Integer Seq., v.11(2008), Article 08.2.8
- [2] . V. Shevelev, *An infinite set of generators of primes based on the Roland idea and conjectures concerning twin primes*, <http://www.arxiv.org/abs/0910.4676> [math. NT].
- [3] . N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences* ([http : //www.research.att.com/~njas](http://www.research.att.com/~njas))